

2004

# Aluminum Maltolate-Induced Toxicity in NT2 Cells Occurs Through Apoptosis and Includes Cytochrome c Release

David A. Dewitt

*Liberty University*, [dadewitt@liberty.edu](mailto:dadewitt@liberty.edu)

Kathleen J. S. Griffioen

Othman Ghribi

Nena Fox

John Savory

Follow this and additional works at: [http://digitalcommons.liberty.edu/bio\\_chem\\_fac\\_pubs](http://digitalcommons.liberty.edu/bio_chem_fac_pubs)

---

## Recommended Citation

Dewitt, David A.; Griffioen, Kathleen J. S.; Ghribi, Othman; Fox, Nena; and Savory, John, "Aluminum Maltolate-Induced Toxicity in NT2 Cells Occurs Through Apoptosis and Includes Cytochrome c Release" (2004). *Faculty Publications and Presentations*. 9. [http://digitalcommons.liberty.edu/bio\\_chem\\_fac\\_pubs/9](http://digitalcommons.liberty.edu/bio_chem_fac_pubs/9)

This Article is brought to you for free and open access by the Department of Biology and Chemistry at DigitalCommons@Liberty University. It has been accepted for inclusion in Faculty Publications and Presentations by an authorized administrator of DigitalCommons@Liberty University. For more information, please contact [scholarlycommunication@liberty.edu](mailto:scholarlycommunication@liberty.edu).

# Aluminum Maltolate-Induced Toxicity in NT2 Cells Occurs Through Apoptosis and Includes Cytochrome *c* Release

Kathleen J.S. Griffioen<sup>1,†</sup>, Othman Ghribi<sup>2</sup>, Nena Fox<sup>3</sup>, John Savory<sup>2</sup>, David A. DeWitt<sup>1,2,\*</sup>

<sup>1</sup>Department of Biology & Chemistry, Liberty University, 1971 University Blvd., Lynchburg, VA 24502 USA

<sup>2</sup>Department of Pathology, University of Virginia, Charlottesville, VA, USA

<sup>3</sup>Department of Microbiology, University of Virginia, Charlottesville, VA, USA

Received 7 October 2003; accepted 17 December 2003

Available online 4 February 2004

## Abstract

Aluminum (Al) compounds are neurotoxic and have been shown to induce experimental neurodegeneration although the mechanism of this effect is unclear. In order to study this neurotoxic effect of Al, we have developed an *in vitro* model system using Al maltolate and human NT2 cells. Al maltolate at 500  $\mu$ M caused significant cell death with a 24-h incubation and this toxicity was even more evident after 48 h. Lower doses of Al maltolate were also effective, but required a longer incubation for cell death. Nuclear fragmentation suggestive of apoptosis was observed as early as three hours and increased substantially through 24 h. Chromatin condensation and nuclear fragmentation were confirmed by electron microscopy. In addition, TUNEL positive nuclei were also observed. The release of cytochrome *c* was demonstrated with Western blot analysis. This *in vitro* model using human cells adds to our understanding of Al neurotoxicity and could provide insight into the neurodegenerative processes in human disease.

© 2004 Elsevier Inc. All rights reserved.

**Keywords:** Aluminum; Apoptosis; Cellular models; Mitochondria; Cytochrome *c*; Neurodegeneration

## INTRODUCTION

Apoptosis or programmed cell death is a regulated process of death in cells that facilitates the removal of extra, aged or damaged cells. In contrast to necrosis, apoptosis is an ordered operation with characteristic apoptotic morphological changes that include nuclear condensation and fragmentation, DNA damage, cell shrinkage, membrane blebbing, and the formation of membrane-bound apoptotic bodies (Huppertz et al., 1999).

Mitochondria are key mediators of apoptosis. Early in apoptosis, the mitochondrial transmembrane

potential ( $\Delta\psi$ ) collapses (Brown et al., 1999), indicating the opening of the permeability transition (PT) pore. When it opens, the PT pore facilitates the equilibration of ions in the intermembrane space, which disconnects the respiratory chain, possibly causing the outer membrane to burst and release cytochrome *c* into the cytosol. Cytochrome *c* release initiates an irreversible cascade that activates programmed cell death (Liu et al., 1996). Several proteins are critical in the regulation of apoptosis. Bax causes the release of cytochrome *c* from mitochondria, driving apoptosis forward (Jürgensmeier et al., 1998). In contrast, Bcl-2 interferes with caspase activity following Bax induction (Rosse et al., 1998), and can also prevent cytochrome *c* from being released at all (Kluck et al., 1997).

Several different triggers have been used to induce apoptosis experimentally including hydrogen peroxide and staurosporine. In addition, aluminum (Al) has been shown to induce death via apoptosis in astrocytes

\* Corresponding author. Tel.: +1-434-582-2228; fax: +1-434-582-2488.

E-mail address: dadewitt@liberty.edu (D.A. DeWitt).

<sup>†</sup> Present address: Institute for Biomedical Sciences, The George Washington University, Washington, DC, USA.

(Suarez-Fernandez et al., 1999).  $\text{AlCl}_3$  has been demonstrated to preferentially accumulate in cultured astrocytic cells (Levesque et al., 2000), resulting in apoptosis. In contrast, Al maltolate accumulates at high levels in neurons, possibly attributable to its lipophilic nature (Levesque et al., 2000).

At 8%, Al is the third most abundant element in the earth's crust. Although it has no known biological function, it is constantly introduced into living systems through soil, water, food, and pharmaceutical agents (Crapper McLachlan, 1986). Despite its apparent lack of positive biological function, it has been reported to be toxic. The first indication of its adverse effects in humans was seen in patients undergoing long term hemodialysis treatment for chronic renal failure. Many of these patients developed hyperaluminemia due to contamination of the dialysis solution with Al and also treatment with oral phosphate binding gels to control hyperphosphatemia. This Al overload resulted in a fatal neurological syndrome (dialysis dementia) and a devastating metabolic bone disease known as dialysis osteodystrophy (reviewed by Mach et al., 1988). Al has been linked to neurotoxicity (Levesque et al., 2000; Perl and Pendlebury, 1986; Roll et al., 1989; Suarez-Fernandez et al., 1999; Troncoso et al., 1985; Tsubouchi et al., 2001) and implicated as a possible causative or contributing factor in neurodegenerative disorders particularly Alzheimer's disease (AD), although the exact mechanism by which it acts is far from clear (Crapper McLachlan, 1986; Shin et al., 1995; Strong, 2001).

Intracisternal administration of Al compounds to New Zealand white rabbits has long been known to induce neurofibrillary pathology (Klatzo et al., 1965). However, apoptosis-related changes and oxidative stress (Ghribi et al., 2001a; Savory et al., 1999) have recently been observed including opening of the permeability transition pore, cytochrome *c* release, Bax and Bcl-2 translocation involving both mitochondria and endoplasmic reticulum, caspase activation, and TUNEL positive nuclei (Ghribi et al., 2001a, 2001d). In these studies, apoptosis was prevented with the administration of cyclosporin A (Ghribi et al., 2001b) and also glial cell derived neurotrophic factor (GDNF) (Ghribi et al., 2001c). The soluble Al compound, Al maltolate, also induces stress to the endoplasmic reticulum when it is administered to rabbits (Ghribi et al., 2001a, 2001d).

Here, we have developed an *in vitro* model of Al toxicity using human NT2 neuronal precursor cells and show that Al maltolate induces apoptosis including the release of cytochrome *c*.

## MATERIALS AND METHODS

### Cell Culture

Human teratocarcinoma (NT2) precursor cells (Stratagene) were grown in DMEM/F-12 (Gibco) growth medium supplemented with 10% (v/v) FBS, 2 mM L-glutamine and 1% (v/v) penicillin–streptomycin and maintained in 5%  $\text{CO}_2$  at 37 °C. For Al treatment, Al maltolate was prepared as a stock solution of 25 mM in sterile water (Berthol et al., 1989) and then passed through 0.2  $\mu\text{m}$  filter. Al was added to growth medium just before use. Unless specified, cells were grown on coverslips in six well plates and allowed to adhere for ~24 h prior to replacement of media containing from 0 to 500  $\mu\text{M}$  Al maltolate.

### Cell Viability (LDH) Assay

Cell viability was assessed using a lactate dehydrogenase assay. NT2 cells were grown in 96 well, round bottom plates at a density of 10,000 cells in 100  $\mu\text{l}$  of media. After Al incubation (5–500  $\mu\text{M}$ ), 50  $\mu\text{l}$  of media was removed and analyzed for LDH activity using the cytotoxicity assay kit from Promega.

### Nuclear and Mitochondrial Visualization

NT2 cells were grown on glass coverslips placed in six-well plates.  $1.5 \times 10^5$  cells in 3 ml media were plated for 24 h prior to addition of Al maltolate. Cells were washed with PBS followed by fixation in 4% formaldehyde for 15 min. Permeabilization was accomplished with an incubation in ethanol:acetic acid (19:1) for 20 min or 0.1% Triton X 100. Nuclei were stained with Hoechst 33258 (Sigma) at 10  $\mu\text{g}/\text{ml}$  for 20 min or VectaShield mounting media with DAPI (Vector Laboratories). Mitochondria were visualized with CMXRos Mitotracker Red (Molecular Probes, Eugene, OR, USA). The mitochondrial polarity-dependent dye was added to culture media at 200 nmol for 20 min before being washed with PBS and fixation in 4% formaldehyde. Coverslips were mounted with VectaShield (Vector Laboratories) and the cells were observed under fluorescence with an Olympus microscope. Pictures were obtained using a digital camera and ImagePro+ software.

### Electron Microscopy

NT2 cells were grown in six-well plates, directly on well bottoms without coverslips. Cells were incubated

in 500  $\mu\text{M}$  Al maltolate for 24 h, rinsed in HBSS and then fixed in gluteraldehyde. Cells were postfixed in 1% osmium tetroxide, embedded and viewed at 150 kV using a Zeiss 902 transmission electron microscope with an EELS spectrometer.

## TUNEL

NT2 cells were grown on glass coverslips in 6-well plates. Media containing 500  $\mu\text{M}$  Al maltolate was added and allowed to incubate at 37 °C at incremental time intervals of 6, and 24 h. Cells were fixed in 4% formaldehyde for 15 min followed by a wash with PBS. DNA damage was assessed using the apoptosis detection kit, fluorescein (Promega) following the manufacturer's instructions.

## Counts

Apoptotic nuclei percentages were determined by obtaining images from 10 random fields on each slide. The total number of cells and the number with apoptotic morphology (strongly condensed or fragmented) were found for each image and then combined to yield a total for each slide. The number of TUNEL positive nuclei was obtained in a similar manner.

## Western Blot

NT2 cells were grown to confluency in T25 flasks. The media was replaced with normal growth media or that containing 500  $\mu\text{M}$  Al maltolate. After six hours in 5%  $\text{CO}_2$  at 37 °C, cells were trypsinized and detached cells were pelleted at 300 g for 5 min. The cell pellet was resuspended in five volumes of homogenizing buffer (20 mM Hepes–KOH (pH 7.5), 10 mM sucrose, 10 mM KCL, 1.5 mM  $\text{MgCl}_2$ , 1 mM EDTA, 1 mM EGTA, 1 mM DTT, 1 mM PMSF, 2 mg/ml aprotinin, 10 mg/ml leupeptin, 5 mg/ml pepstatin and 12.5 mg/ml of *N*-acetyl-Leu-Leu-Norleu-Al) (Liu et al., 1996) and homogenized in a Teflon homogenizer. The resulting suspension was centrifuged for 5 min in a micro-centrifuge. The supernatant was then centrifuged at 100,000  $\times$  g using a Beckman Optima LE-80K ultracentrifuge for 30 min to obtain cytosolic extract. The protein concentration was determined using the Bradford method and BSA as a standard (BioRad). Cytosolic and mitochondrial extract was mixed with 2x Lamelli sample buffer and frozen at –80 °C until used for electrophoresis. 10  $\mu\text{g}$  of protein was loaded onto precast 4–20% gradient SDS-PAGE gels (BioRad) and subsequently transferred to PVDF membrane.

Membranes were blocked with 5% non-fat dry milk and incubated with monoclonal antibody to human cytochrome *c* (Pharmingen, Lexington, KY, USA). A monoclonal antibody to  $\beta$ -actin (Sigma) was used as a gel loading control. Protein bands were visualized with chemiluminescence (Vector Laboratories). An antibody to cytochrome oxidase IV (Molecular Probes) was used to determine the extent of mitochondrial contamination of the cytosol.

## RESULTS

### Aluminum is a Potent Neurotoxin for Cultured NT2 Cells

Cell viability was determined by measuring lactate dehydrogenase (LDH) release into tissue culture media. This assay allowed determination of the appropriate cell density and Al maltolate concentrations for further study. 500  $\mu\text{M}$  and 250  $\mu\text{M}$  Al maltolate-induced noticeable death of NT2 cells after a 24-h incubation (Fig. 1) and effectively depleted cells from the culture after 48 h. Lower doses of Al maltolate (50–100  $\mu\text{M}$ ) were also neurotoxic, but required longer incubation times to elicit similar neuronal death observed at 250  $\mu\text{M}$  and 500  $\mu\text{M}$  at 24 h.

### Aluminum-Induced Neurotoxicity Occurs via Apoptosis

To assess the type of death evoked by Al maltolate, we used Hoechst 33258 or DAPI to examine nuclear morphology. Nuclear fragmentation consistent with apoptotic morphology was observed as early as three hours following incubation with Al maltolate. At 24 h, numerous fragmented and condensed nuclei could clearly be observed in the culture (Fig. 2B). The percentage of apoptotic nuclei had increased at 6 h and rose dramatically through 24 h (Fig. 2C). Under phase microscopy, the size of some cells with apoptotic nuclei was clearly reduced (data not shown).

Chromatin condensation and nuclear fragmentation were further confirmed by electron microscopy. Nuclei of control cultures appeared healthy and round, fully enclosed within the nuclear membrane (Fig. 3A). Al-treated cultures contained nuclei that appeared irregular and separated into fragments (Fig. 3B). In addition, nuclei in experimental cultures contained patches of condensed DNA consistent with apoptotic




Fig. 1. Al maltolate induces cell death Cell viability was determined by measuring the amount of LDH released into the culture media at 2 h (A); 24 h (B) and 48 h (C). Incubation with 250–500  $\mu$ M Al maltolate-induced substantial cell death at 24 and 48 h. Lower doses of Al maltolate also induced death of NT2 cells, however, longer incubation times were required. Data is representative of three independent experiments with similar trends. Values are  $\pm$ S.D.

morphology. Many of the separated nuclear fragments were surrounded by membrane although some of the nuclear vesicles appeared ruptured.

Al-treated cells exhibited DNA damage, a key marker of apoptosis, as indicated by TUNEL positive nuclei (Fig. 4D). A significant increase in the number of cells with DNA damage and nuclear fragmentation was observed at six hours and a 10-fold increase in damage was observed by 24 h (Fig. 4F). This occurred with a clear decrease in cell number in Al maltolate-treated cultures (Fig. 4E).




Fig. 2. DAPI staining of NT2 cells revealed large round or oval nuclei (A). Following 24 h incubation in Al maltolate, the nuclei take on a bright, condensed (arrowhead) or fragmented (arrows) morphology (B). The percentage of nuclei with a fragmented, apoptotic morphology had increased as early as 6 h and increased further at 24 h (C). Values are means  $\pm$  S.E.M. and representative from three independent experiments. Scale bar = 40  $\mu$ m ( $*P < 0.001$ ).

### Al-treated NT2 Cells Undergo Apoptosis Including the Release of Cytochrome c

CMXRos is a mitochondrial specific dye that is taken up in a membrane polarity dependent manner.




Fig. 3. Electron microscopy confirmed nuclear condensation in Al maltolate-treated cultures. Nuclei of cells in control cultures (A) appeared healthy with a round, even morphology. In contrast, nuclei of Al maltolate-treated cultures (B) were typically fragmented (arrows).

Therefore, the dye can be used to demonstrate mitochondria with an intact membrane potential. Polarized mitochondria were observed in nearly all of the cells even following treatment with Al maltolate (Fig. 4C).

Release of cytochrome *c* to the cytoplasm was confirmed with Western blotting. Cytosolic extract was isolated from control and Al maltolate-treated cultures after a 6 h incubation. Cytochrome *c* was detected in the cytoplasm of Al maltolate-treated cells to a much greater extent than controls (Fig. 5). In order to verify that the cytochrome *c* in the cytosolic fraction was not due to mitochondrial contamination, an antibody to cytochrome oxidase IV was used. While present in the mitochondrial fraction, cytochrome oxidase IV was absent from the cytosolic fraction.

## DISCUSSION

Although the neurotoxic effects of Al have been well established in conditions such as dialysis encephalopathy (Mach et al., 1988), the mechanism by which it elicits its effects remain to be elucidated. Here, we show that Al maltolate toxicity in human NT2 cells occurs via apoptosis and includes cytochrome *c* release. Our results indicate this neurotoxicity is time and concentration dependent, and occurs via apoptosis as shown by nuclear fragmentation and chromatin condensation, DNA damage, and cytochrome *c* release.

Al maltolate-induced substantial cell death of the NT2 precursor cells with a 24-h incubation. Concentrations of 250–500  $\mu\text{M}$  Al maltolate were highly toxic, while lower concentrations required longer incubations to cause significant death. Similar concentrations of Al have been used in vitro and shown to be neurotoxic (Levesque et al., 2000; Suarez-Fernandez et al., 1999; Tsubouchi et al., 2001). In particular, Al has a narrow effective range with a large difference in outcome (Roll et al., 1989). Pulse or low concentration exposure appears to be less toxic than higher concentrations (Kashiwagi et al., 1998). We found that lower concentrations of Al maltolate (50–100  $\mu\text{M}$ ) did induce death but required longer incubations for a significant effect. It is possible that Al must first accumulate intracellularly to a critical level before triggering cell death. Such accumulation is likely to be concentration and time dependent as well as impacted by the Al species administered (Levesque et al., 2000). Accumulation below a certain threshold may have little or no effect. Alternatively, sub-lethal accumulated concentrations insufficient to induce apoptosis may still negatively impact the cytoskeleton or intracellular transport.

Cytochrome *c* release is a key event in the initiation of apoptosis (Liu et al., 1996). This release generally coincides with opening of the mitochondrial PT pore and loss of mitochondrial membrane polarity. An earlier study reported that cytochrome *c* could be released by staurosporine without concomitant depolarization of the mitochondrial membrane potential (Krohn et al., 1999). Al-induced apoptosis appears to occur through a similar mechanism. Previously, Al was shown to induce mitochondrial permeability transition (Toninello et al., 2000). However, when Al was bound to the PT pore, it held it in an intermediate, partially open position. It is possible that this is sufficient to trigger cytochrome *c* release.

While a causal link between Al and Alzheimer's disease has not been demonstrated, several epidemiological studies have shown a statistically significant




Fig. 4. Numerous NT2 cells can be observed following a 24 h incubation in control media (CMXRos, red A). Very few of these cells were TUNEL positive (B, green). However, after 24 h in Al maltolate (C) a clear reduction in cell number has occurred. Many of these cells were TUNEL positive (D, green). The number of cells in control cultures increases over 24 h but decreases in Al maltolate-treated cultures (E) which has more TUNEL positive nuclei as well (F). Values are means  $\pm$  S.E.M. and representative from three independent experiments with similar trends (\* $P < 0.5$ , \*\* $P < 0.001$ ). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)




Fig. 5. Western blot analysis revealed a greater amount of cytochrome *c* in the cytoplasm of Al maltolate-treated cultures (Al) compared with controls (Ctl). The absence of cytochrome oxidase IV in the cytoplasm indicated that mitochondrial contamination of the cytoplasm was negligible.

positive link (reviewed by Flatten, 2001). Nonetheless, aluminum induces many of the same biochemical and pathological changes that occur in AD (reviewed by Strong). Regardless of any lack of evidence for a causative link, aluminum has potential as a trigger to cause cellular changes that mimic aspects of AD pathology (Perl and Pendlebury, 1986; Huang et al., 1997). If similar pathways are activated, Al-induced pathology may prove useful as a model to understand the sequence of events that occurs as neurons die in AD brain. New Zealand white rabbits have been used successfully in this regard (Ghribi et al., 2001a, 2001b, 2001d; Kowall et al., 1989; Klatzo et al., 1965; Munoz-Garcia et al., 1986; Savory et al., 1999, 2001).

Aside from inducing apoptosis, Al negatively affects cytoskeletal proteins and axonal transport (Kashiwagi et al., 1998; Shea et al., 1997). Indeed, pulse exposure to Al maltolate blocks fast axonal transport leading to the accumulation of NF-L proteins in rat cortical neurons (Kashiwagi et al., 1998). Disrupted transport has been shown in AD (Cash et al., 2003; Praprotnik et al., 1996; Richard et al., 1989; Terry, 1996) and also amyotrophic lateral sclerosis (Williamson and Cleveland, 1999), however relatively few studies have examined this aspect of the disease (Kasa et al., 2000). Nonetheless, numerous factors implicated in AD including oxidative stress (de la Monte et al., 2000; Perry and Smith, 1997; Smith et al., 1995), APO  $\epsilon$  (Tesseur et al., 2000), and APP-L (Torroja et al., 1999) have all been shown to affect axonal transport. The observation that A $\beta$ PP can serve as a kinesin cargo receptor (Kamal et al., 2000, 2001) as well as the inhibition of kinesin-dependent transport by tau overexpression (Ebnet et al., 1998; Stamer et al., 2002) should not be overlooked. It would be interesting to see whether tau overexpression and Al affect intracellular transport via a similar mechanism.

Apoptosis has been suggested to play a role in AD pathology (Cotman and Anderson, 1995; Su et al., 1997) however, either the full cascade is not activated or appears to be aborted (Perry et al., 1998a, 1998b; Raina et al., 2001). An intriguing possibility is that apoptosis-related changes, oxidative stress, and axonal transport disruption are linked in a way that promotes neurodegeneration and involves cytoskeletal disruption (Srivastava et al., 1998). Since Al contributes to all of these factors (Ghribi et al., 2001b; Kashiwagi et al., 1998; Pratico et al., 2002), the model presented here may prove useful in understanding the mechanisms leading to neuronal death in neurodegenerative diseases.

## ACKNOWLEDGEMENTS

This work was funded by the Jeffress Memorial Trust, the National Institutes of Health (AG-020996), an undergraduate research grant from the Virginia Academy of Sciences (K.J.S.G.) and an undergraduate research grant from the Liberty University Biology Alumni. The authors thank M.S. Forbes, Ph.D. for assistance with electron microscopy and P. Pramoonjago, Ph.D. for assistance with western blotting.

## REFERENCES

- Berthol RL, Herman MM, Savory J, Carpenter RM, Sturgill BC, Katsetos CD, Vandenberg SR, Wills MR. A long-term intravenous model of aluminum maltol toxicity in rabbits: tissue distribution, hepatic, renal, and neuronal cytoskeletal changes associated with systemic exposure. *Toxicol Appl Pharmacol* 1989;98(1):58–74.
- Brown GC, Nicholls DG, Cooper CE, editors. *Mitochondria and cell death*. New Jersey: Princeton University Press, 1999.
- Cash AD, Aliev G, Siedlak SL, Nunomura A, Fujioka H, Zhu X et al. Microtubule reduction in Alzheimer's disease and aging is independent of tau filament formation. *Am J Pathol* 2003; 162:1623–7.
- Cotman CW, Anderson AJ. A potential role for apoptosis in neurodegeneration and Alzheimer's disease. *Mol Neurobiol* 1995;10(1):19–45.
- Crapper McLachlan DR. Aluminum and Alzheimer's disease. *Neurobiol Aging* 1986;7:525–32.
- de la Monte SM, Neely TR, Cannon J, Wands JR. Oxidative stress and hypoxia-like injury cause Alzheimer-type molecular abnormalities in central nervous system neurons. *Cell Mol Life Sci* 2000;57(10):1471–81.
- Ebnet A, Godemann R, Stamer K, Illenberger S, Trinczek B, Mandelkow E. Overexpression of tau protein inhibits kinesin-dependent trafficking of vesicles, mitochondria and endoplasmic reticulum: implications for Alzheimer's disease. *J Cell Biol* 1998;143(3):777–94.



- Flatten TP. Aluminum as a risk factor in Alzheimer's disease, with emphasis on drinking water. *Brain Res Bull* 2001;55(2):187–96.
- Ghribi O, DeWitt DA, Forbes MS, Herman MM, Savory J. Co-involvement of mitochondria and endoplasmic reticulum in regulation of apoptosis: changes in cytochrome *c*, Bcl-2 and Bax in hippocampus of aluminum-treated rabbits. *Brain Res* 2001a;903:66–73.
- Ghribi O, DeWitt DA, Forbes MS, Arad A, Herman MM, Savory J. Opening of the mitochondrial permeability transition pore and cytochrome *c* release is an early response to neuronal injury in aged rabbits. *J Alzheimer Dis* 2001b;3(4):387–91.
- Ghribi O, Herman MM, Forbes MS, DeWitt DA, Savory J. GDNF protects against aluminum-induced apoptosis in rabbits by upregulating Bcl-2 and Bcl-X<sub>L</sub>, and inhibiting mitochondrial Bax translocation. *Neurobiol Dis* 2001c;8:764–73.
- Ghribi O, Herman MM, DeWitt DA, Forbes MS, Savory J. A $\beta$ (1-42) and aluminum induce stress in the endoplasmic reticulum in rabbit hippocampus, involving nuclear translocation of GADD 153 and NF- $\kappa$ B. *Mol Brain Res* 2001d;96:30–8.
- Huang Y, Herman MM, Liu J, Katssetos CD, Wills MR, Savory J. Neurofibrillary lesions in experimental aluminum-induced encephalopathy and Alzheimer's disease share immunoreactivity for amyloid precursor protein, A $\beta$ ,  $\alpha$ <sub>1</sub>-antichymotrypsin and ubiquitin-protein conjugates. *Brain Res* 1997;771:213–20.
- Huppertz B, Frank HG, Kaufmann P. The apoptosis cascade—morphological and immunohistochemical methods for its visualization. *Anat Embryol* 1999;200:1–18.
- Jürgensmeier JM, Xie Z, Deveraux Q, Ellerby L, Bredesen D, Reed JC. Bax directly induces release of cytochrome *c* from isolated mitochondria. *Proc Nat Acad Sci USA* 1998;95:4997–5002.
- Kamal A, Stokin GB, Yang Z, Xia CH, Goldstein LS. Axonal transport of amyloid precursor protein is mediated by direct binding to the kinesin light chain subunit of kinesin-I. *Neuron* 2000;28(2):449–59.
- Kamal A, Almenar-Queralt A, LeBlanc JF, Roberts EA, Goldstein LS. Kinesin-mediated axonal transport of a membrane compartment containing beta secretase and presenilin-1 requires APP. *Nature* 2001;414:643–8.
- Kasa P, Papp H, Kovacs I, Forgon M, Penke B, Yamaguchi H. Human amyloid- $\beta$  1-42 applied in vivo inhibits the fast axonal transport of proteins in the sciatic nerve of rat. *Neurosci Lett* 2000;278:117–9.
- Kashiwagi Y, Nakamura Y, Miyamae Y, Hashimoto R, Takeda M. Pulse exposure of cultured rat neurons to aluminum maltol affected the axonal transport system. *Neurosci Lett* 1998;252:5–8.
- Klatzo I, Wisniewski HM, Streicher E. Experimental production of neurofibrillary degeneration. Part 1. Light microscopic observation. *J Neuropathol Exp Neurol* 1965;24:187–99.
- Kluck RM, Bossy-Wetzel E, Green DR, Newmeyer DD. The release of cytochrome *c* from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science* 1997;275:1132–6.
- Kowall NW, Pendlebury WW, Kessler JB, Perl DP, Beal MF. Aluminum-induced neurofibrillary degeneration affects a subset of neurons in rabbit cerebral cortex, basal forebrain and upper brainstem. *Neuroscience* 1989;29(2):329–37.
- Krohn AJ, Wahlbrink T, Prehn JHM. Mitochondrial depolarization is not required for neuronal apoptosis. *J Neurosci* 1999;19(17):7394–404.
- Levesque L, Mizzen CA, McLachlan DR, Fraser PE. Ligand specific effects on aluminum incorporation and toxicity in neurons and astrocytes. *Brain Res* 2000;877:191–202.
- Liu X, Kim CN, Yang J, Jemmerson R, Wang X. Induction of Apoptotic Program in Cell-Free Extracts: Requirement for dATP and Cytochrome *c*. *Cell* 1996;86:147–57.
- Mach JR, Korchik WP, Mahowald MW. Dialysis dementia. *Clin Geriatr Med* 1988;4:853–67.
- Munoz-Garcia D, Pendlebury WW, Kessler JB, Perl DP. An immunocytochemical comparison of cytoskeletal proteins in aluminum-induced and Alzheimer-type neurofibrillary tangles. *Acta Neuropathol (Berl)* 1986;70(3/4):243–8.
- Perl DP, Pendlebury WW. Aluminum neurotoxicity—potential role in the pathogenesis of neurofibrillary tangle formation. *Can J Neurol Sci* 1986;13(4, suppl):441–5.
- Perry G, Nunomura A, Smith MA. A suicide note from Alzheimer disease neurons? *Nat Med* 1998;4(8):897–8.
- Perry G, Nunomura A, Lucassen P, Lassmann H, Smith MA. Apoptosis and Alzheimer's disease. *Science* 1998;282:1268–9.
- Perry G, Smith MA. A central role for oxidative damage in the pathogenesis and therapeutics of Alzheimer's disease. *Alzheimer Dis* 1997;2(7):319–24.
- Praprotnik D, Smith MA, Richey PL, Vinters HV, Perry G. Filament heterogeneity within the dystrophic neurites of senile plaques suggests blockage of fast axonal transport in Alzheimer's disease. *Acta Neuropathol* 1996;91:226–35.
- Pratico D, Uryu K, Sung S, Tang S, Trojanowski JQ, Lee VM. Aluminum modulates brain amyloidosis through oxidative stress in APP transgenic mice. *FASEB J* 2002;16(9):1138–40.
- Raina AK, Hochman A, Zhu X, Rottkamp CA, Nunomura A, Siedlak SL, Boux H, Castellani RJ, Perry G, Smith M. Abortive apoptosis in Alzheimer's disease. *Acta Neuropathol (Berl)* 2001;101(4):305–10.
- Richard S, Brion P, Couck AM, Flament-Durand J. Accumulation of smooth endoplasmic reticulum in Alzheimer's disease: New morphological evidence of axoplasmic flow disturbances. *J Submicrosc Cytol Pathol* 1989;21(3):461–7.
- Roll M, Banin E, Meiri H. Differentiated neuroblastoma cells are more susceptible to aluminum toxicity than developing cells. *Toxicology* 1989;63:231–7.
- Rosse T, Olivier R, Monney L, Ragre M, Conus S, Fellay I, Jansen B, Borner C. Bcl-2 prolongs cell survival after Bax-induced release of cytochrome *c*. *Nature* 1998;391:496–9.
- Savory J, Rao JKS, Letada P, Herman MM. Age-related hippocampal changes in Bcl-2:Bax ratio, oxidative stress, redox-active iron and apoptosis associated with aluminum-induced neurodegeneration: increased susceptibility with aging. *NeuroToxicology* 1999;20:805–18.
- Savory J, Ghribi O, Forbes MS, Herman MM. The rabbit model system for studies of aluminum-induced neurofibrillary degeneration: relevance to human neurodegenerative disorders. Aluminum and Alzheimer's disease: the science that describes the link. Elsevier Science, 2001. p. 203–19.
- Shea TB, Wheeler E, Jung C. Aluminum inhibits neurofilament assembly, cytoskeletal incorporation, and axonal transport. Nature of aluminum-induced perikaryal neurofilament accumulations as revealed by subunit turnover. *Mol Chem Neuropathol* 1997;32:17–39.
- Shin RW, Lee VMY, Trojanowski JQ. Neurofibrillary pathology and aluminum in Alzheimer's disease. *Histol Histopathol* 1995;10:969–78.

- Smith MA, Sayre LM, Monnier VM, Perry G. Radical ageing in Alzheimer's disease. *Trends Neurosci* 1995;18:172–6.
- Srivastava RK, Srivastava AR, Korsmeyer SJ, Nesterova M, Cho-Chung YS, Longo DL. Involvement of microtubules in the regulation of Bcl-2 phosphorylation and apoptosis through cyclic AMP-dependent protein kinase. *Mol Cell Biol* 1998;18(6):3509–17.
- Stamer K, Vogel R, Thies E, Mandelkow E, Mandelkow EM. Tau blocks traffic of organelles, neurofilaments, and APP vesicles in neurons and enhances oxidative stress. *J Cell Biol* 2002;156:1051–63.
- Strong MJ. Aluminum as an experimental neurotoxicant: the neuropathology and neurochemistry. In Exley C, editor, *Aluminum and Alzheimer's disease: the science that describes the link*. Elsevier Science, 2001.
- Su JH, Deng G, Cotman CW. Bax protein expression is increased in Alzheimer's brain: correlations with DNA damage, Bcl-2 expression, and brain pathology. *J Neuropathol Exp Neurol* 1997;56(1):86–93.
- Suarez-Fernandez MB, Soldado AB, Sanz-Mendel A, Vega JA, Novelli A, Fernandez-Sanchez MT. Aluminum-induced degeneration of astrocytes occurs via apoptosis and results in neuronal death. *Brain Res* 1999;835:125–36.
- Terry RD. The pathogenesis of Alzheimer disease: an alternative to the amyloid hypothesis. *J Neuropathol Exp Neurol* 1996;55(10):1023–5.
- Tesseur I, Van Dorpe J, Bruynseels K, Bronfman F, Sciot R, Van Lommel A, Van Leuven F. Prominent axon and disruption of axonal transport in transgenic mice expressing human apolipoprotein E4 in neurons of brain and spinal cord. *Am J Pathol* 2000;157(5):1495–510.
- Torroja L, Chu H, Kotovsky I, White K. Neuronal overexpression of APPL, the *Drosophila* homologue of the amyloid precursor protein (APP), disrupts axonal transport. *Curr Biol* 1999;9(9):489–92.
- Troncoso JC, Hoffman PN, Griffin JW, Hess-Kozlow KM, Price DL. Aluminum intoxication: a disorder of neurofilament transport in motor neurons. *Brain Res* 1985;342(1):172–5.
- Tsubouchi R, Htay HH, Murakami K, Haneda M, Yoshino M. Aluminum-induced apoptosis in PC12D cells. *Biometals* 2001;14(2):181–5.
- Toninello A, Clari G, Marcon M, Tognon G, Zatta P. Aluminum as an inducer of the mitochondrial permeability transition. *J Biol Inorg Chem* 2000;5:612–23.
- Williamson TL, Cleveland DW. Slowing of axonal transport is a very early event in the toxicity of ALS-linked SOD1 mutants to motor neurons. *Nature Neurosci* 1999;2(1):50–6.